

Ncstn-Flox

系統名	C57BL/6Smoc- <i>Ncstn</i> ^{em1(flox)Smoc}
SMOC番号	NM-CKO-241011
維持形態	Developing

遺伝子の概要

Gene Symbol Ncstn	Synonyms	Nct; Aph2; AA727311; D1Dau13e; Kiaa0253; mKIAA0253; 9430068N19Rik
	NCBI ID	59287
	MGI ID	1891700
	Ensembl ID	ENSMUSG00000003458
	Human Ortholog	NCSTN

説明

These strains carry loxP sites flanking exon 3 of Ncstn gene. When crossed with a Cre recombinase-expressing strain, this strain is useful in eliminating tissue-specific conditional expression of Ncstn gene.

*Literature published using this strain should indicate: Ncstn-Flox mice (Cat. NO. NM-CKO-241011) were purchased from Shanghai Model Organisms Center, Inc..

病気の予測

Chronic Myeloid Leukemia	表現型	MGI:5009037 Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with VAV1-cre mice.
	参考文献	Klinakis A, Lobry C, Abdel-Wahab O, Oh P, Haeno H, Buonamici S, van De Walle I, Cathelin S, Trimarchi T, Araldi E, Liu C, Ibrahim S, Beran M, Zavadil J, Efstratiadis A, Taghon T, Michor F, Levine RL, Aifantis I, A novel tumour-suppressor function for the Notch pathway in myeloid leukaemia. <i>Nature</i> . 2011 May 12;473(7346):230-3

Urinary Bladder Cancer	表現型	MGI:5779426 Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Upk2-cre mice.
	参考文献	Rampias T, Vgenopoulou P, Avgeris M, Polyzos A, Stravodimos K, Valavanis C, Scorilas A, Klinakis A, A new tumor suppressor role for the Notch pathway in bladder cancer. <i>Nat Med.</i> 2014 Oct;20(10):1199-205
Schizophrenia	表現型	MGI:5800496 Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Olig1-Cre mice.
	参考文献	Dries DR, Zhu Y, Brooks MM, Forero DA, Adachi M, Cenik B, West JM, Han YH, Yu C, Arbellal J, Nordin A, Adolfsson R, Del-Favero J, Lu QR, Callaerts P, Birnbaum SG, Yu G, Loss of Nicastin from Oligodendrocytes Results in Hypomyelination and Schizophrenia with Compulsive Behavior. <i>J Biol Chem.</i> 2016 May 27;291(22):11647-56
Chronic Myeloid Leukemia	表現型	MGI:5009036 Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Mx1-cre mice.
	参考文献	Klinakis A, Lobry C, Abdel-Wahab O, Oh P, Haeno H, Buonamici S, van De Walle I, Cathelin S, Trimarchi T, Araldi E, Liu C, Ibrahim S, Beran M, Zavadil J, Efstratiadis A, Taghon T, Michor F, Levine RL, Aifantis I, A novel tumour-suppressor function for the Notch pathway in myeloid leukaemia. <i>Nature.</i> 2011 May 12;473(7346):230-3

表現型データ

No data