

Myl1-NLS-Cre

系統名 C57BL/6Smoc-*Myl1*^{em1(NLS-iCre-pA)Smoc}

SMOC番号 NM-KI-200124

維持形態 Repository Live

遺伝子の概要

Gene Symbol Myl1	Synonyms	Mylf; MLC1f; MLC3f; AI325107
	NCBI ID	<u>17901</u>
	MGI ID	<u>97269</u>
	Ensembl ID	ENSMUSG00000061816
	Human Ortholog	MYL1

説明

A NLS-iCre-pA expression cassette was knocked into the Myl1 gene start codon site via CRISPR/Cas9 mediated recombination.

応用分野: Cre-mediated recombination can be detected in the skeletal muscle of Myl1-Cre mice. When crossed with a strain carrying a gene flanked by loxP sites, the flanked gene will be removed in cells expressing cre.

*Literature published using this strain should indicate: Myl1-NLS-Cre mice (Cat. NO. NM-KI-200124) were purchased from Shanghai Model Organisms Center, Inc..

表現型デロタ






Fig. 1 Cre-mediated recombination in the skeletal muscle of Myl1^{Cre/+}; Rosa26^{tdTomato/+} mouse. TdTomato(red) expression can be detected in the skeletal muscle of Myl1^{Cre/+}; Rosa26^{tdTomato/+} mouse.




Fig. 2 Detection of tdTomato(red) in various tissues of Myl1^{Cre/+}; Rosa26^{tdTomato/+} mice. Cre mediated recombination can be detected in the skeletal muscle. Tdtomato expression can also be detected in individual cells derived from liver, myocardium and brown fat. Tdtomato expression can not be observed in the brain and stomach. (For more detailed information please contact our technical advisor.)