

Nkx2-5-IRES-Cre

系統名 C57BL/6Smoc-*Nkx2-5*^{em1(IRES-Cre-WPRE-polyA)Smoc}

SMOC番号 NM-KI-200152

維持形態 Sperm cryopreservation

遺伝子の概要

Gene Symbol Nkx2-5	Synonyms	Csx; Nkx2.5; tinman; Nkx-2.5
	NCBI ID	<u>18091</u>
	MGI ID	<u>97350</u>
	Ensembl ID	ENSMUSG00000015579
	Human Ortholog	NKX2-5

説明

A IRES-Cre-WPRE-polyA expression cassette was knocked into the Nkx2-5 gene stop codon site. **応用分野**: Cre recombinase tool; Nkx2.5 acts as a key transcription factor and plays essential roles for heart formation. By mating the reporter mice with Cre-expressing mice, Nkx2-5 positive cells derived from double-positive mice were permanently labeled by fluorescent protein. When crossed with mice carrying a targeted floxed alleles, this strain is useful in eliminating tissue-specific conditional expression of the gene.

*Literature published using this strain should indicate: Nkx2-5-IRES-Cre mice (Cat. NO. NM-KI-200152) were purchased from Shanghai Model Organisms Center, Inc..

表現型デロタ






Fig. 1 Cre-mediated recombination in the heart of Nkx2- $5^{\text{Cre/+}}$; Rosa2 $6^{\text{tdTomato/+}}$ mouse. TdTomato(red) expression can be detected in the myocardial cells and vascular cells of Nkx2- $5^{\text{Cre/+}}$; Rosa2 $6^{\text{tdTomato/+}}$ mouse.




Fig. 2 Detection of tdTomato(red) in various tissues of Nkx2-5^{cre/+}; Rosa26^{tdTomato/+} mice. Cre mediated recombination can be detected in the heart, bronchus, pulmonary alveoli and submandibular gland. Tdtomato expression can be also detected in individual cells derived from the brain, retina, skin, kidney, pancreas, intestine, colon, spleen and thymus, except for pituitary and muscle. (For more detailed information please contact our technical advisor.)

