

# Braf-Flox

系統名	C57BL/6Smoc- <i>Braf</i> <sup>em1(flox)Smoc</sup>
SMOC番号	NM-CKO-200073
維持形態	Repository Live

## 遺伝子の概要

<b>Gene Symbol</b> <b>Braf</b>	<b>Synonyms</b>	B-raf; Braf2; Braf-2; C87398; AA120551; AA387315; AA473386; C230098H17; D6Ert631e; 9930012E13Rik
	<b>NCBI ID</b>	<a href="#">109880</a>
	<b>MGI ID</b>	<a href="#">88190</a>
	<b>Ensembl ID</b>	<a href="#">ENSMUSG00000002413</a>
	<b>Human Ortholog</b>	BRAF

## 説明

These mice carry loxP sites flanking exon 5 of Braf gene. When crossed with a Cre recombinase-expressing strain, this strain is useful in eliminating tissue-specific conditional expression of Braf gene.

**応用分野:** Study on the correlation between meiosis and RET signal transduction in oocytes

\*Literature published using this strain should indicate: Braf-Flox mice (Cat. NO. NM-CKO-200073) were purchased from Shanghai Model Organisms Center, Inc..

## 病気の予測

<b>Melanoma</b>	<b>表現型</b>	<a href="#">MGI:3843341</a> Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Tyr-cre/ERT2 mice.
	<b>参考文献</b>	Dhomen N, Reis-Filho JS, da Rocha Dias S, Hayward R, Savage K, Delmas V, Larue L, Pritchard C, Marais R, Oncogenic Braf induces melanocyte senescence and melanoma in mice. <i>Cancer Cell</i> . 2009 Apr 7;15(4):294-303

<b>Thyroid Gland Papillary Carcinoma</b>	<b>表現型</b>	<a href="#">MGI:5779643</a> Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with TPO-cre mice.
	<b>参考文献</b>	Franco AT, Malaguarnera R, Refetoff S, Liao XH, Lundsmith E, Kimura S, Pritchard C, Marais R, Davies TF, Weinstein LS, Chen M, Rosen N, Ghossein R, Knauf JA, Fagin JA, Thyrotrophin receptor signaling dependence of Braf-induced thyroid tumor initiation in mice. Proc Natl Acad Sci U S A. 2011 Jan 25;108(4):1615-20
<b>Thyroid Gland Papillary Carcinoma</b>	<b>表現型</b>	<a href="#">MGI:5780077</a> Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Tg-cre/ERT2 mice.
	<b>参考文献</b>	Charles RP, Iezza G, Amendola E, Dankort D, McMahon M, Mutationally Activated BRAFV600E Elicits Papillary Thyroid Cancer in the Adult Mouse. Cancer Res. 2011 Jun 1;71(11):3863-71
<b>Skin Melanoma</b>	<b>表現型</b>	<a href="#">MGI:5902132</a> Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Pten-Flox(NM-CKO-18004) and Tyr-cre/ERT2 mice.
	<b>参考文献</b>	Dankort D, Curley DP, Cartlidge RA, Nelson B, Karnezis AN, Damsky WE Jr, You MJ, DePinho RA, McMahon M, Bosenberg M, Braf(V600E) cooperates with Pten loss to induce metastatic melanoma. Nat Genet. 2009 May;41(5):544-52
<b>Prostate Cancer</b>	<b>表現型</b>	<a href="#">MGI:5543907</a> Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Pten-Flox(NM-CKO-18004) and Nkx3-1-Cre mice.
	<b>参考文献</b>	Wang J, Kobayashi T, Floc'h N, Kinkade CW, Aytes A, Dankort D, Lefebvre C, Mitrofanova A, Cardiff RD, McMahon M, Califano A, Shen MM, Abate-Shen C, B-Raf activation cooperates with PTEN loss to drive c-Myc expression in advanced prostate cancer. Cancer Res. 2012 Sep 15;72(18):4765-76

<b>Langerhans-Cell Histiocytosis</b>	<p><b>表現型</b></p> <p><a href="#">MGI:6192274</a></p> <p>Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with Cd207-Cre mice.</p> <p><b>参考文献</b></p> <p>Berres ML, Lim KP, Peters T, Price J, Takizawa H, Salmon H, Idoyaga J, Ruzo A, Lupo PJ, Hicks MJ, Shih A, Simko SJ, Abhyankar H, Chakraborty R, Leboeuf M, Beltrao M, Lira SA, Heym KM, Bigley V, Collin M, Manz MG, McClain K, Merad M, Allen CE, BRAF-V600E expression in precursor versus differentiated dendritic cells defines clinically distinct LCH risk groups. <i>J Exp Med.</i> 2014 Apr 7;211(4):669-83</p>
<b>Cardiofaciocutaneous Syndrome</b>	<p><b>表現型</b></p> <p><a href="#">MGI:5902221</a></p> <p>Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with CAG-cre mice.</p> <p><b>参考文献</b></p> <p>Inoue S, Moriya M, Watanabe Y, Miyagawa-Tomita S, Niihori T, Oba D, Ono M, Kure S, Ogura T, Matsubara Y, Aoki Y, New BRAF knockin mice provide a pathogenetic mechanism of developmental defects and a therapeutic approach in cardio-facio-cutaneous syndrome. <i>Hum Mol Genet.</i> 2014 Dec 15;23(24):6553-66</p>
<b>cardiofaciocutaneous syndrome</b>	<p><b>表現型</b></p> <p><a href="#">MGI:6164161</a></p> <p>Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with CAG-cre mice.</p> <p><b>参考文献</b></p> <p>Moriya M, Inoue S, Miyagawa-Tomita S, Nakashima Y, Oba D, Niihori T, Hashi M, Ohnishi H, Kure S, Matsubara Y, Aoki Y, Adult mice expressing a Braf Q241R mutation on an ICR/CD-1 background exhibit a cardio-facio-cutaneous syndrome phenotype. <i>Hum Mol Genet.</i> 2015 Dec 20;24(25):7349-60</p>

<b>Langerhans-Cell Histiocytosis</b>	<b>表現型</b>	<p><a href="#">MGI:6192275</a> Note: The expected phenotype(s) may be observed in the above-mentioned mice that bred with CAG-cre mice.</p>
	<b>参考文献</b>	<p>Berres ML, Lim KP, Peters T, Price J, Takizawa H, Salmon H, Idoyaga J, Ruzo A, Lupo PJ, Hicks MJ, Shih A, Simko SJ, Abhyankar H, Chakraborty R, Leboeuf M, Beltrao M, Lira SA, Heym KM, Bigley V, Collin M, Manz MG, McClain K, Merad M, Allen CE, BRAF-V600E expression in precursor versus differentiated dendritic cells defines clinically distinct LCH risk groups. <i>J Exp Med.</i> 2014 Apr 7;211(4):669-83</p>

## 表現型データ

No data