

Anxa10-2A-CreERT2

系統名	C57BL/6Smoc- <i>Anxa10</i> ^{em1(2A-CreERT2-Wpre-pA)Smoc}
SMOC番号	NM-KI-200312
維持形態	Repository Live

遺伝子の概要

Gene Symbol Anxa10	Synonyms
	NCBI ID 26359
	MGI ID 1347090
	Ensembl ID ENSMUSG00000031635
	Human Ortholog ANXA10

説明

2A-CreERT2-Wpre-pA expression cassette was knocked into the Anxa10 gene stop codon site.

応用分野: These mice express tamoxifen induced cre recombinase from the Anxa10 locus. By mating the reporter mice with CreERT2-expressing mice, reporter gene expression can be detected in gastric mucosa epithelial cells after tamoxifen treatment. And the mice are useful for generating conditional mutations in gastric mucosa epithelial cells. This strain may useful for in the research of gastric carcinoma.

*Literature published using this strain should indicate: Anxa10-2A-CreERT2 mice (Cat. NO. NM-KI-200312) were purchased from Shanghai Model Organisms Center, Inc..

表現型データ




Fig.1 CreERT2-mediated recombination in the stomach of Anxa10^{CreERT2/+}; Rosa26^{tdTomato/+} mouse.

TdTomato(red) expression can be detected in the gastric mucosal epithelial cell of Anxa10^{CreERT2/+}; Rosa26^{tdTomato/+} mouse after tamoxifen treatment.




Fig.2 Detection of tdTomato (red) in various tissues of Anxa10^{CreERT2/+}; Rosa26^{tdTomato/+} mice after tamoxifen treatment.

CreERT2-mediated recombination in the gastric mucosal cells and hair follicle cells can be induced by tamoxifen. A small degree of leakiness were detected in the thymus prior to tamoxifen exposure. Besides, tdtomato expression can not be detected in the brain, eyes, heart, lung, liver, submandibular gland, kidney, ovary, uterus, intestine, colon, spleen, brown fat, and

muscle. (For more detailed information please contact our technical advisor.)
