

# Il4-Luc-2A-EGFP

系統名	C57BL/6Smoc- <i>Il4</i> <sup>em1(Luc-eGFP)Smoc</sup>
SMOC番号	NM-KI-00021
維持形態	Repository Live

## 遺伝子の概要

<b>Gene Symbol</b> <b>IL4</b>	<b>Synonyms</b>	IL-4; BSF-1
	<b>NCBI ID</b>	<a href="#">16189</a>
	<b>MGI ID</b>	<a href="#">96556</a>
	<b>Ensembl ID</b>	<a href="#">ENSMUSG00000000869</a>
	<b>Human Ortholog</b>	IL4
<b>Gene Symbol</b> <b>Il4</b>	<b>Synonyms</b>	BSF-1; IL-4
	<b>NCBI ID</b>	<a href="#">16189</a>
	<b>MGI ID</b>	<a href="#">96556</a>
	<b>Ensembl ID</b>	<a href="#">ENSMUSG00000000869</a>
	<b>Human Ortholog</b>	IL4

## 説明

A Luciferase-2A-EGFP-pA cassette was inserted after ATG the start codon. The intrinsic promoter drives the expression of reporter genes, which reflects the activity of Il4.

**応用分野:** Inflammation and immunity, gene tracing

\*Literature published using this strain should indicate: Il4-Luc-2A-EGFP mice (Cat. NO. NM-KI-00021) were purchased from Shanghai Model Organisms Center, Inc..

## 表現型データ




Fig1. Time course of mean fluorescence intensity on the ventral of Il4-Luc-2A-EGFP mice after 12 mg/mL Con.A administration.

Abbr. HO, homozygous; HE, heterozygous; WT, wild type.




Fig2. Individual fluorescence intensity on the ventral (A) and dorsal (B) of Il4-Luc-2A-EGFP mice treated with Con.A.